Figure 1. The beautiful but dangerous *Podostroma cornu-damae*, courtesy Taylor Lockwood. Recent poisonings in Japan were apparntely caused by confusion with the similarly colored club mushroom *Clavulinopsis miyabeana*.



## (that you've never heard of)

## Stephen F. Nelsen

Department of Chemistry, University of Wisconsin, 1101 University Avenue, Madison, WI 53706-1396 USA

rofessor Hideo Tomioka, a friend of mine recently sent word from Nagoya, Japan, of a dangerous mushroom that has been making headlines there. According to the August 26, 2011 edition of Yomiuri Shinbun, Japan's largest daily, a mushroom known as kaen-dake (in Romanji) has been expanding its range in Shiga Prefecture. Alarms have been raised as this region is adjacent to the huge Chūkyō metropolitan area which is the 3rd largest in Japan (8.74 million people) and includes the city of Nagoya. The newspaper article included a photo of the bright red coral like fungus Podostroma cornu-damae (Fig. 1) and cautioned readers to be very careful of this mushroom. The fungus was unfamiliar to both of us, but there is an uncommon (and not known to be dangerous) species of Podostroma in North America (Podostroma alutaceum, Fig. 2 and 3).

Several poisonings have been reported in Japan, resulting from consumption of Podostroma cornu-damae. In 1999, one of a group of five people from Niigata prefecture died two days after consuming about 1 gram (0.035 oz) of a fruit body that had been soaked in sake. In 2000, an individual from Gunma prefecture died after eating the fried mushroom. Symptoms associated with consumption in these cases included stomach pains, changes in perception, decrease in the number of leukocytes and thrombocytes, peeling skin on the face, hair loss, and shrinking of the cerebellum, resulting in speech impediment and problems with voluntary movement (Saikawa et al., 2001). In another instance, an autopsy revealed multiple organ failure, including acute kidney failure, liver necrosis and disseminated intravascular coagulation (Koichi et al., 2003). In one case of poisoning, the patient suffered from hemophagocytosis, in addition to severe leukocytopenia and thrombocytopenia seven days after ingesting the fungus. Plasmapheresis and administration of granulocyte colony-stimulating factor

were used to treat the blood disorders. The authors suggested that these treatments, in addition to the large volume of administered intravenous fluid—9 liters (2.4 US gal) over a 12-hour period—were responsible for his successful recovery (Suzuki et al., 2002).

The poisoning symptoms are similar to

We are certainly lucky that this exceptionally poisonous fungus does not occur here. Professor Tomioka wrote that simply touching it is said to lead to the swelling of your flesh!

Podostroma cornu-damae is an ascomycete fungus, a member of the order Hypocreales (Pezizomycotina), and previously has gone by other names. According to Index Fungorum the fungus was originally described as *Hypocrea cornu-damae* by Patoullard (1895) and was found "in lignis putidus in Thibet, orient." *Podostroma cornu-damae* is next attributed to Boedijn (1934) (and see Doi, 1973). It was published in Saccardo's famous Sylloge Fungorum (1905), which was a comprehensive list of all of the names that had been used for mushrooms and other fungi (and was the only such list of its kind).



Figure 2. The North American species, *Podostroma alutaceum*, found in Iowa. Courtesy of Jim Frink and the Prairie States Mushroom Club.

those observed previously with animals that have consumed trichothecene mycotoxins. Japanese researchers detected the presence of the macrocyclic trichothecenes: satratoxin H, satratoxin H 12', 13'-diacetate, satratoxin H 12'-acetate, and satratoxin H 13'-acetate. When grown in liquid culture the fungus additionally produces roridin E, verrucarin J, and satratoxin H. With the exception of verrucarin J, a 0.5 gram (0.018 oz) dose of all of these compounds, when injected into the abdomen of mice, will result in their death the following day (Saikawa et al., 2001).

## **References Cited**

Boedijn, K.B. 1934. The genus *Podostroma* in The Netherlands Indies. *Bulletin du Jardin Botanique Buitenzorg* 13: 269–275.

Doi, Y. 1973. Revision of the Hypocreales with cultural observations. *V. Podostroma* giganteum Imai, *P. cornu-damae* (Pat.) Boedijn and *Hypocrea pseudogelatinosa* sp. nov. *Report of the Tottori Mycological Institute* 10: 421–427.

Koichi, M., T. Haruo, Y. Toshihiro, O. Masami, N. Sadao, and K. Koichiro. 2003. Case report: food poisoning to death by *Podostroma cornu-damae*, its case history and autopsy findings. (In Japanese) *Acta Criminologiae et Medicinae Legalis Japonica* 69(1): 14–20.

Patouillard, N.T. 1895. Enumeration des champignons recoltes par les RR. PP. Farges et Soulie, dans le Thibet oriental et le Sutchuen. (In French) *Bulletin de la Société Mycologique de France* 11: 196–199.

Saccardo P.A., and D. Saccardo. 1905. Supplementum universale. Pars VI. Hymenomycetae-Laboulbeniomycetae. (In Latin) *Sylloge Fungorum* 17: 799.

Saikawa, Y., H. Okamoto, T. Inui, M. Makabe, T. Okuno, T. Suda, K. Hashimoto, and M. Nakata. 2001. Toxic principles of a poisonous mushroom *Podostroma cornudamae*. *Tetrahedron* 57(39): 8277–8281.

Suzuki M, Y. Katoh, H. Kumagai, M. Saitoh, H. Ishikawa, H. Itoh, and K. Shimazu K. 2002. Successful treatment in a case of *Podostroma cornu-damae* poisoning, a deadly poisonous mushroom. (In Japanese) *Chudoku Kenkyu* 15(2): 177–182.



Figure 3. The North American species, *Podostroma alutaceum*, found in Iowa and courtesy of Jim Frink and the Prairie States Mushroom Club.

Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund Fund
Send me a full, one-year subscription to FUNGI.
I want to receive <b>5</b> big, colorful issues!
(4 seasonal issues + annual special issue)
🔲 \$38 USA 🛄 \$40 Canada or Mexico 🛄 \$50 for all other Intl. addresses
Name
Address
City State Zip
Country Email
Place CREDIT CARD orders at the FUNGI website,
www.fungimag.com
or drop your check in the mail today! Make check out to "FUNGI" and send it to: FUNGI & P.O. Box 8 & 1925 Hwy. 175 & Richfield, Wisconsin 53076-0008 USA
If you give us your email address, we'll send you subscription reminders, announcements, and notices of information placed on the website. Your email address will NOT be traded or sold, and will not be shared with anyone not directly affiliated with FUNGI.

FUNGI (ISSN 1941-4943) is published five times per year (four seasonal issues plus a special issue) by FUNGI, P.O. Box 8, 1925 Hwy. 175, Richfield, Wisconsin 53076-0008, USA. Subscriptions are \$38 per year for USA residents; \$40 for residents of Canada and Mexico; \$50 for all others. Checks should be made out to FUNGI. For credit card orders please see our Web site: www.fungimag.com

PUBLISHER'S NOTES: Although many wild mushrooms are quite palatable, some are deadly poisonous. It is advisable to avoid eating any wild organisms, including fungi, unless absolutely certain of identification. And although some mushroom species are edible for many people, those same species may cause allergic reactions or illness in others. When in doubt, throw it out. FUNGI wants to ensure that all readers are long-term subscribers. It is a good idea to have any wild mushroom checked by an expert before eating them. It should be understood that the Publisher and all Editors are not responsible for any consequences of ingesting wild mushrooms. Furthermore, the Publisher and all Editors are not engaged, herein, in the rendering of any medical advice or services. All readers should verify all information and data before administering any drug, therapy, or treatment discussed herein. Neither the Editors nor the Publisher accepts any responsibility for the accuracy of the information or consequences from the use or misuse of the information contained herein. Unauthorized reproduction of published content of FUNGI is strictly forbidden, and permission for reproduction must be obtained by application in writing to the Publisher.

COPYRIGHT ©2011 by FUNGI.

All rights reserved. Printed in the U.S.A.